


PDL1-Multiple Drugs and Multiple Assays in Development Drug and IVD Company Perspectives



Debra J Rasmussen

Senior Director Global Regulatory Affairs

Janssen Pharmaceuticals

October 8, 2015



Janssen is proud to feature artwork created by people affected by the illnesses and diseases we are committed to treating and preventing.

OVERVIEW

- Definitions
- History of Immuno-Oncology
- The Case of PD-L1
- Companion Diagnostic Development

US FDA Definition of IVDs



IVDs are a subset of medical devices which are

“ . . . **reagents, instruments, and systems** intended for use in the **diagnosis of disease or other conditions**, including a determination of the state of health, in order to cure, **mitigate, treat, or prevent disease or its sequelae.**”

Such products are intended for use in the collection, preparation, and examination of specimens taken from the human


(21 CFR 809.3)



PHARMACEUTICAL COMPANIES
OF **Johnson & Johnson**

In Vitro Diagnostic (IVD)

IVD: Reagents and systems used for testing specimens taken from the body




Comprised of many unique technology segments



Regulatory: What is being Approved?

IVD comprises reagents and systems used for testing specimens taken from the body, typically performed in a lab
FDA reviews 3 components of the test and their interactions



Companion Diagnostic

- “An IVD companion diagnostic device is an in vitro diagnostic device that provides information that is essential for the safe and effective use of a corresponding therapeutic product.” *
- In most circumstances (new drug or new indication), an IVD companion diagnostic device and its corresponding therapeutic product should be **approved or cleared contemporaneously** for the use indicated in the therapeutic product labeling.

**FDA definition.*



PHARMACEUTICAL COMPANIES
OF 

| 10/5/2015 | 6



U.S. Food and Drug Administration
Protecting and Promoting Public Health

www.fda.gov

Current Companion Diagnostic Examples

Many successful companion diagnostic/therapeutic co-approvals

– www.fda.gov/companiondiagnostics

CoDx Complexities - Easy Cases:

- One drug, one disease indication, one test, one allele:
e.g., Abbott VYSIS ALK Break Apart FISH Probe Kit for Xalkori® (crizotinib)
- One test, one indication, more than one drug, same alleles
➢ E.g., QIAGEN *therascreen* KRAS RGQ PCR Kit– for CRC for two therapeutics Erbitux® (cetuximab) and Vectibix® (panitumumab).
- Originally one drug, one indication:
➢ HER2 and Herceptin –breast cancer (Tests - Dako IHC / Vysis PathVysion FISH)
➢ Follow on tests demonstrated method comparison, development of CAP guidelines
➢ Expanded the indication to gastric cancer
➢ Other drugs for the same analyte - Perjeta, Kadcyca



PHARMACEUTICAL COMPANIES
OF Johnson & Johnson

| 7 |



U.S. Food and Drug Administration
Protecting and Promoting Public Health

www.fda.gov

CoDx Complexities

One indication, More than one drug, Two tests- Same gene but different allele representation.

- BRAFV600 mutation:
 - Roche cobas BRAF V600 Mutations Test for Zelboraf® (vemurafenib)
 - BioMérieux THxID BRAF Kit for Tafinlar® (dabrafenib) and Mekinist® (trametinib)
- EGFR Activating Mutations
 - Roche cobas EGFR Mutation Test for Tarceva® (erlotinib)
 - Qiagen *therascreen* EGFR RGQ PCR Kit for Gilotrif® (afatinib)





PHARMACEUTICAL COMPANIES
OF 

Cancer immunotherapy
(immuno-oncology) is the use of the immune system to treat cancer.

The immune system is the body's natural defense system.


- Cancer cells can be very different(foreign) from normal cells so the immune system attacks them.
- Cancer cells often find ways to disguise themselves as normal cells or, similar to viruses, cancer cells can change over time (mutate) and therefore escape from the immune response.
- Natural immune response to cancer cells is often not strong enough to fight off cancer cells.
- Immuno-oncology works by activating our immune system to recognize cancer cells and destroy them.






The Case of PD-L1

a Innate immune resistance



b Adaptive immune resistance



Nature Reviews | Cancer



U.S. Food and Drug Administration
Protecting and Promoting Public Health

www.fda.gov

PD-L1 IHC Methods in Development

	Hopkins	BMS	Merck	Genentech
mAb clone	5H1	28-8	22C3	SP142
Automated	No	Yes	Yes	Yes
Staining location scored	Membrane	Membrane	Membrane	Membrane
Cell type(s) scored	Tumor cells	Tumor cells	Tumor and/or infiltrating immune cells	Infiltrating immune cells
Positive cutoff	≥ 5%	≥ 5%	≥ 1%	≥1% to ≥10% ("IHC 1-2-3")

Topalian 2015

➤ **Note:** *These assays are evolving, pending further clinical correlations*



PHARMACEUTICAL COMPANIES
OF *Johnson & Johnson*

13

PD-1 Blocking Antibodies

- **Nivolumab (ONO-4538, BMS-936558, or MDX1106)**, developed by Ono Pharmaceutica and Medarex (Medarex was later acquired by **Bristol-Myers Squibb**) and marketed as **Opdivo**, is a fully humanized IgG4 monoclonal antibody that blocks PD-1
- **Pembrolizumab** (formerly **MK-3475** and **lambrolizumab**, trade name **Keytruda; Merck**) is a humanized monoclonal IgG-4k antibody that blocks PD-1

Anti-PD-L1 Antibodies

- **MPDL3280A (Roche)** is a humanized IgG-1k monoclonal anti-PD-L1 antibody. It is genetically engineered to modify the Fc domain, thereby impairing the antibody-dependent cellular cytotoxicity of PD-L1 expressing cells
- **MEDI4736 (AstraZeneca)** is a humanized IgG-1k monoclonal antibody that blocks PD-L1

Nivolumab (Opdivo/BMS)

- **FDA approval:** Treated Squamous cell non-small cell lung cancer (NSCLC) that has progressed based on CheckMate-017, A Phase III study of Nivolumab vs Docetaxel in 2nd Line Squamous Cell. *Responses were observed independent of PD-L1 status*
- “Importantly, PD-LI emerged as a clear predictive factor for the benefit of nivolumab,” Dr. Paz-Ares stated at the ASCO 2015 press conference
- Recent: Checkmate 057, a Phase III, open-label, randomized study of Nivolumab versus docetaxel in previously treated patients with advanced or metastatic non-squamous NSCLC

Pembrolizumab (Keytruda/Merck)




Figure 3. Progression-free Survival.

Shown are Kaplan–Meier estimates of progression-free survival according to the proportion score (PS) — the percentage of neoplastic cells with membranous PD-L1 staining — for 356 patients in the training and validation groups who had slides that were sectioned within 6 months before staining (Panel A), including 294 previously treated patients (Panel B) and 62 previously untreated patients (Panel C).


“These data in combination with responses among patients with a proportion score of less than 1% suggest that tumor PD-L1 expression is not associated with the ideal test characteristics of approved genetically based biomarkers. It is likely that tumor PD-L1 expression alone does not accurately assess the “dynamic immune microenvironment.”

Garon et al. Pembrolizumab for the Treatment of Non–Small-Cell Lung Cancer NEJM 2015

MPDL3280A (Roche)

- FDA breakthrough designation: treatment of patients with PD-L1–positive non–small cell lung cancer (NSCLC) that has progressed during or after platinum-based chemotherapy, as well as a targeted therapy for patients with EGFR- or ALK-positive tumors. (Phase I study; PDL1 as inclusion criterium)
- Ongoing: -A Phase 2 Study of MPDL3280A Compared With Docetaxel in Patients With Locally Advanced or Metastatic NSCLC Who Have Failed Platinum Therapy "POPLAR". Interim results: MPDL3280A reduced the risk of death by 53% ([OS]; [HR]=0.47) in people whose cancer expressed the highest levels of PD-L1 compared with docetaxel. An improvement in OS was also observed in people who had medium and high (HR=0.56) or any level of PD-L1 expression (HR=0.63), as characterized by a test being developed by Roche.

MPDL3280A (Roche)



The association of response to MPDL3280A treatment and tumour-infiltrating immune cell PD-L1 expression reached statistical significance (NSCLC, $P=0.015$), while the association with tumour cell PD-L1 expression did not (NSCLC, $P=0.920$)

Figure 3 | Antitumour activity of MPDL3280A by immunohistochemistry (IHC) tumour-infiltrating immune cell (IC) and biomarker status. a, Table

Herbst et al "Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients" Nat Let 2015



U.S. Food and Drug Administration
Protecting and Promoting Public Health

www.fda.gov

Problematic for the following reasons

- High potential for mismatched approved drug/device combination in the clinical setting. Patient treatment may not be based on testing with the matched CoDx.
- FDA approvals are for a specific drug/CoDx combination, so the labeling applies only to that particular combination. Performance across different clinical decision points (cut-offs) may not be established.
- Laboratories are not able to assess the impact of discordance between tests in the absence of clinical outcome data.
- Laboratories are not expected to have more than one assay/platform to detect the same analyte.



PHARMACEUTICAL COMPANIES
OF Johnson & Johnson

19

THANK YOU!





U.S. Food and Drug Administration
Protecting and Promoting Public Health

www.fda.gov

Candidate CoDx Complexities – The Case of PD-L1

- 4-8 drugs in development
- Parallel development programs
- Various trial designs
- Multiple anti-PD-L1 immunohistochemistry (IHC) companion diagnostics
 - Different test for each drug



PHARMACEUTICAL COMPANIES
OF *Johnson & Johnson*

| 21 |